



## A (ii) RETURNING HOME FOLLOWING HOSPITAL TREATMENT FOR FRACTURED HIP (IMPROVEMENT)

Percentage of patients discharged back to usual place of residence within 28 days of emergency admission to hospital with a hip fracture, aged 65 and over (age and sex standardised), 1999/00 to 2000/01




**A (ii) RETURNING HOME FOLLOWING HOSPITAL TREATMENT FOR FRACTURED HIP (IMPROVEMENT)**

Percentage of patients discharged back to usual place of residence within 28 days of emergency admission to hospital with a hip fracture, aged 65 and over (age and sex standardised), 1999/00 to 2000/01



## A (ii) RETURNING HOME FOLLOWING HOSPITAL TREATMENT FOR FRACTURED HIP (IMPROVEMENT)

Percentage of patients discharged back to usual place of residence within 28 days of emergency admission to hospital with a hip fracture, aged 65 and over (age and sex standardised), 1999/00 to 2000/01



## A (ii) RETURNING HOME FOLLOWING HOSPITAL TREATMENT FOR FRACTURED HIP (IMPROVEMENT)

Percentage of patients discharged back to usual place of residence within 28 days of emergency admission to hospital with a hip fracture, aged 65 and over (age and sex standardised), 1999/00 to 2000/01

